

Population Ecology of Sacramento Splittail


Fred Feyrer¹, Jim Hobbs², Shawn Acuna³, Brian Mahardja⁴,
Lenny Grimaldo⁵, Melinda Baerwald², Rachel Johnson⁶, & Swee Teh²

*¹US Geological Survey, ²UC Davis, ³Metropolitan Water District of Southern California, ⁴California Department of Water Resources,
⁵ICF International, & ⁶National Marine Fisheries Service*



Sacramento Splittail *Pogonichthys macrolepidotus*



- Large migratory cyprinid with broad environmental tolerances
- Endemic to the San Francisco Estuary and tributaries
- Only surviving member of genus
- Formerly listed as threatened by US Fish and Wildlife Service





Young-of-year






Adult






Listed
1999






Petaluma-Napa
Population




Central Valley
Population




Petaluma-Napa
Population

Central Valley
Population

Estuary ?





Petaluma-Napa
Population

The diagram shows a map of California with three highlighted regions: a red oval in the northwestern coastal area labeled 'Petaluma-Napa Population', a green circle in the central coastal area labeled 'Estuary ?', and a large blue oval covering the central valley area labeled 'Central Valley Population'. A black silhouette of the state of California is shown in the lower-left corner, with the highlighted regions overlaid on it.

Central Valley
Population

Estuary ?


Metapopulation Criteria

- (1) Inhabit discrete breeding habitat patches.
- (2) Connected via dispersal.
- (3) Traits not synchronous.

Key Research Questions

- (i) Are individuals from the two populations found disproportionately in specific geographic regions or habitats of the estuary?
- (ii) Do individuals within aggregations/schools originate from the same population?
- (iii) Do individuals within a population or within a geographic region exhibit differences in demographic or health metrics relative to others?

Splittail Distribution




Splittail Distribution




Splittail Habitat Affinities


Central Valley subpopulation GAM: 80% deviance explained




Petaluma/Napa subpopulation GAM: 61% deviance explained




Splittail Population Distribution




Splittail Age and Growth



Splittail Growth Rate




Splittail Health and Condition



Splittail Population Connectivity

Genetic Assignment


- Unassigned
- Petaluma-Napa population
- Central Valley population



2010

Dry Winter/Spring

Central Valley population **not observed** in Petaluma-Napa breeding habitat




2011


Wet Winter/Spring

Central Valley population **observed** in Petaluma-Napa breeding habitat


Splittail Population Connectivity



Splittail Population Connectivity



Metapopulation structure



Conclusions

- (i) Individuals from the two populations **ARE NOT** found disproportionately in specific geographic regions or habitats of the estuary.

Conclusions

- (i) Individuals from the two populations **ARE NOT** found disproportionately in specific geographic regions or habitats of the estuary.
- (ii) Individuals within aggregations/schools **DO** originate from the same population.

Conclusions

- (i) Individuals from the two populations **ARE NOT** found disproportionately in specific geographic regions or habitats of the estuary.
- (ii) Individuals within aggregations/schools **DO** originate from the same population.
- (iii) Individuals within a population **SOMETIMES** or within a geographic region **YES** exhibit differences in demographic or health metrics relative to others.

Conclusions

- (i) Individuals from the two populations **ARE NOT** found disproportionately in specific geographic regions or habitats of the estuary.
- (ii) Individuals within aggregations/schools **DO** originate from the same population.
- (iii) Individuals within a population **SOMETIMES** or within a geographic region **YES** exhibit differences in demographic or health metrics relative to others.
- (iv) Island-mainland metapopulation structure.

Conclusions

- (i) Individuals from the two populations **ARE NOT** found disproportionately in specific geographic regions or habitats of the estuary.
- (ii) Individuals within aggregations/schools **DO** originate from the same population.
- (iii) Individuals within a population **SOMETIMES** or within a geographic region **YES** exhibit differences in demographic or health metrics relative to others.
- (iv) Island-mainland metapopulation structure.
- (v) Climate variability drives connectivity and therefore possibly gene flow.

Conclusions

- (i) Individuals from the two populations **ARE NOT** found disproportionately in specific geographic regions or habitats of the estuary.
- (ii) Individuals within aggregations/schools **DO** originate from the same population.
- (iii) Individuals within a population **SOMETIMES** or within a geographic region **YES** exhibit differences in demographic or health metrics relative to others.
- (iv) Island-mainland metapopulation structure.
- (v) Climate variability drives connectivity and therefore possibly gene flow.
- (vi) Implications for management in a changing climate and landscape.

Connecting Fish Tissue Selenium Concentrations to Sources and
Exposure in a Dynamic Estuary: The Case of Sacramento Splittail
Robin Stewart, USGS

Thursday 3:40
Room 308-310



Acknowledgements

**Marty Gingras
Ramiro Soto
Eric Santos
Nick Sakata
Nick van Ark
Colin Grant
Doug Obegi**

**Randy Baxter
Bill Harrell
Anke Mueller-Solger
Matt Young
Andrew Bibian
Louise Conrad
Brian Schreier**

**Ted Sommer
Kevin Reece
Michael MacWilliams
Larry Brown
Nann Fangue
Bernie May
Karen Gehrts**

**Erwin van Nieuwenhuyse
Scott Waller
Ted Foin
Matt Nobriga
Dan Riordon
Amy Chandros
Bobby Coalter**

**U.S. Bureau of Reclamation
Delta Science Program
California Department of Fish and Wildlife
California Department of Water Resources**